

- Abbott, M.B., J.C. Bathurst, J.A. Cunge, P.E. O’Connell and J. Rasmussen. 1986. An introduction to the European Hydrological System – Systeme Hydrologique Europeen “SHE” 2: Structure of a physically based distributed modeling system. *Journal of Hydrology* 87: 61-77.
- Arnold, J.G., R. Srinivasan, R.S. Muttiah, and J.R. Williams. 1998. Large area hydrologic modeling and assessment. Part I: Model development. *J. American Water Resources Association*, 34(1): 73-89; Adapted for this study by NASA Ames Research Center and California State University Monterey Bay.
- Ault, T., R. Velzeboer, and R. Zammit. 2000. Influence of nutrient availability on phytoplankton growth and community structure in the Port Adelaide River, Australia: bioassay assessment of potential nutrient limitation. *Hydrobiologia* 429: 89-103.
- Bredrossian, T.L. and Custis, K. 2002. Memorandum to Review of July 2002 EPA Analysis of Impacts of Timberland Management on Water Quality. Department of Conservation, California Geological Survey.
- Bellows B. 2001. Nutrient Cycling in Pastures. *Livestock System Guide*. <http://www.attra.ncat.org>.
- Bicknell, B.R., J.C. Imhoff, J.L. Kittle, A.S. Donigian, and R.C. Johanson. 1993. Hydrological Simulation Program–Fortran (HSPF): User’s manual for release 10.0. EPA 600/3-84-0466. U.S. Environmental Protection Agency, Environmental Research Laboratory, Athens, GA.
- Biggs, B.J.F. 1988. Algal proliferations in New Zealand’s shallow stony foothills-fed rivers: toward a predictive model. *Verh. Int. Ver. Limnol.* 23:1405–11.
- Blake, M.C., Jr., R. W. Groyer, and R.E. Stramski. 2002. Geologic map and map database of western Sonoma, northernmost Marin, and southernmost Mendocino counties, California. . Pamphlet to accompany miscellaneous field studies map MF-2402, v.1.0. United States Geological Survey (USGS).
- Blus, L.J., and C.J. Henny. 1997. Field studies on pesticides and birds: Unexpected unique relations. *Ecological Applications* 7: 1125-1132.
- Borah, D.K., M. Bera. 2004. Watershed-scale hydrologic and nonpoint-source pollution models: Review of applications. *Transactions of the ASAE* 47: 789-803.

- California Data Exchange Center (CDEC). April 10, 2007. <http://wdl.water.ca.gov/gw>.
- California Department of Water Resources (DWR). December 1975. Evaluation of ground water resources in Sonoma County. Volume 1: geologic and hydrologic data. DWR Bulletin 118-4.
- California Department of Water Resources (DWR). 1982. Evaluation of ground water resources in Sonoma County. Volume 2: Santa Rosa Plain. DWR Bulletin 118-4.
- California Department of Water Resources (DWR). October 2003. California's Groundwater Bulletin 118-Update.
- Cardwell, G.T. 1958. Geology and ground water in the Santa Rosa and Petaluma Valley areas, Sonoma County, California. USGS Water Supply Paper 1427.
- Carnell, R.E., and C.A. Senior. 1998. Changes in mid-latitude variability due to greenhouse gases and sulphate aerosols. *Climate Dynamics* 14:369–383.
- Carr, G., and P. A. Chambers. 1998. Macrophyte growth and sediment phosphorous and nitrogen in a Canadian prairie river. *Freshwater Biology* 39: 525-536.
- Cassel, E.A., R.L. Kort, D.W. Meals, S.G. Aschmann, J.M. Dorioz, and D.P. Anderson. 2001. Dyanmic phosphorus mass balance modeling of large watersheds: long-term implications of management strategies. *Water Science and Technology* 43: 153-162.
- CDFG. 1995a. Fish Species of Special Concern in California, Hardhead. California Department of Fish and Game, Habitat Conservation Planning Branch. http://www.CDFG.ca.gov/hcpb/cgi-bin/more_info.asp?idKey=ssc_tespp&specy=fish&query=Mylopharodon%20conocephalus.
- CDFG. 1995b. Fish Species of Special Concern in California, Tule Perch. California Department of Fish and Game, Habitat Conservation Planning Branch. http://www.CDFG.ca.gov/hcpb/cgi-bin/read_one.asp?specy=fish&idNum=38.
- CDFG. 2000a. California Department of Fish and Game stream inventory report: North Fork Santa Rosa Creek. Revised April 14, 1006.
- CDFG. 2000b. California Department of Fish and Game stream inventory report: Santa Rosa Creek. Revised April 14, 1006.
- CDFG. 2005a. California Department of Fish and Game stream inventory report: Copeland Creek Revised April 14, 1006.
- CDFG. 2005b. California Department of Fish and Game stream inventory report: Blucher Creek. Revised April 14, 1006.
- CH2M Hill. 1989. Santa Rosa subregional water reclamation system, long-term detailed studies, Phase I: Development of reclamation alternative. Volume 2: Receiving water evaluation-Laguna de Santa Rosa. Prepared for the City of Santa Rosa.

- Chen, C.W., J. Herr, and L. Weintraub. 2000. Watershed analysis risk management framework (WARMF): Update One – A decision support system for watershed analysis and total maximum daily load calculation, allocation and implementation. Publication No. 1005181, Electric Power Research Institute, Palo Alto, CA. 2001.
- Chen, C.W., J.W. Herr, R.A. Goldstein, G. Ice, T. Cundy. 2005. Retrospective comparison of watershed analysis risk management framework and Hydrologic Simulation Program–Fortran applications to Mica Creek watershed. *Journal of Environmental Engineering*, ASCE 131: 1277- 1284.
- Chorley, R.J. and B.A. Kennedy. 1971. *Physical geography: A system approach*. London. Prentice-Hall.
- City of Santa Rosa. January 1992. Final environmental impact report, Santa Rosa Creek master plan.
- City of Santa Rosa. 2005. 2004-05 Annual report for the Santa Rosa area storm water permit 2, term 2, part V.
- Coey, R, S. Nossaman-Pearce, C. Brooks, and Z. Young. 2002 (Draft). California Department of Fish and Game Russian River basin fisheries restoration plan. California Department of Fish and Game.
- Cook, D., and J. Martini-Lamb. 2001. Copeland Creek restoration project monitoring plan. http://www.scwa.ca.gov/_pdf/copeland_creek_restoration.pdf
- Cook, D., and D. Manning. 2002. Data report 1999-2001: Russian River Basin steelhead and coho salmon monitoring program pilot study. http://www.scwa.ca.gov/_pdf/russian_river_steelhead_coho.pdf
- Craig, D. and P.L. Williams. 1998. “Willow Flycatcher (*Empidonax traillii*)” in *The Riparian Bird Conservation Plan: a strategy for reversing the decline of riparian-associated birds in California*. California Partners in Flight. http://www.prbo.org/calpif/htmldocs/species/riparian/willow_flycatcher.htm
- Crooks, J.A. 2002. Characterizing ecosystem-level consequences of biological invasions: the role of ecosystem engineers. *Oikos* 97: 153-166.
- Crowder, A., and D.S. Painter, 1991. Submerged macrophytes in Lake Ontario: Current knowledge, importance, threats to stability, and needed studies. *Canadian Journal of Fisheries and Aquatic Sciences*, 48: 1539-1545.
- Custer, C.M. 1993. “Life history, traits and habitat needs of the redhead” in *Waterfowl Management Handbook*. LaCrosse, WI: U.S. Fish and Wildlife Service http://www.nwrc.usgs.gov/wdb/pub/wmh/13_1_11.pdf
- Davis, W.S. and T.P. Simons, eds. 1995. *Biological assessment and criteria: Tools for resource planning and decision making*. Lewis Publishers. Boca Raton, FL.

- Decker, K. 2007. The fate of nitrogen and phosphorus in relation to inundation of the Laguna de Santa Rosa floodplain. A white paper for conceptual model development. (Available through Chris Potter, NASA Ames Research Center, cpotter@mail.arc.nasa.gov.)
- Delta Regional Ecosystem Restoration Implementation Plan (DRERIP). May 2005. Framework for the development of DRERIP ecosystem conceptual models. Department of Conservation, Division of Mines and Geology, Special Report 120.
- Diaz, M.M. and F.L. Pedrozo. 1996. Nutrient limitation in Andean-Patagonian lakes at latitude 40-41°S. *Archiv für Hydrobiologie* 138: 123-143.
- Dodds, W.K., V.H. Smith, and K. Lohman. 2002. Nitrogen and phosphorus relationships to benthic algal biomass in temperate streams. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 865-874.
- Downing, J.A., C.W. Osenberg, and O. Sarnelle. 1999. Metanalysis of marine nutrient enrichment experiments: variation in the magnitude of nutrient limitation. *Ecology* 80: 1157-1167.
- Duever, M.J. 2005. Big Cypress regional ecosystem conceptual ecological model. *Wetlands* 25: 843-853.
- Dwernychuk, L.W. and D.A. Boag. 1973. Effect of herbicide-induced changes in vegetation on nesting ducks. *Canadian Field Naturalist* 87:155-165.
- Federal Emergency Management Agency (FEMA). 1997. Flood insurance study, Sonoma County, unincorporated areas. Volume 3 of 4.
- Fenn, M.E., J.S. Baron, E.B. Allen, H.M. Rueth, K.R. Nydick, L. Geiser, W.D. Bowman, J.O. Sickman, T. Meixner, and D.W. Johnson. 2003. Ecological effects of nitrogen deposition in the western United States. *BioScience* 53:404-420.
- Fitz, H.C., E.B. DeBellevue, R. Costanza, R. Boumans, T. Maxwell, L. Wainger, F.H. Sklar. 1996. Development of a general ecosystem model for a range of scales and ecosystems. *Ecological Modeling* 88: 263-295.
- Flosi, G., S. Downie, J. Hopelain, M. Bird, R. Coey, and B. Collins. 1998. California salmonid stream habitat restoration manual, 3rd edition. California Department of Fish and Game, Sacramento, CA.
- Fredrickson, L.H., and F.A. Reid. 1986. "Wetland and riparian habitats: a nongame management overview", 59-96 in J.B. Hale, L.B. Best, and R.L. Clawson, eds; *Management of nongame wildlife in the Midwest: a developing art. Proceedings of the 47th Midwest Fish and Wildlife Conference. The Wildlife Society, North Central Section, Grand Rapids, Michigan.*
- Giorgi, F., L. Mearns, S. Shields, and L. McDaniel. 1998. Regional nested model simulations of present day and future climate over the central Great Plains of the United States. *Climatic Change* 40:457-493.

- Gluesenkamp, D., and J. Wirka. 2006. Sonoma Valley vernal pools: Is nitrogen pollution harming fragile ecosystems? *The Ardeid*: 8-10.
- Gratwicke, B., and B.E. Marshall. 2001. The impact of *Azolla filiculoides* Lam. on animal biodiversity in streams in Zimbabwe. *African Journal of Ecology* 39: 216–218.
- Greenway, M. 1997. Nutrient content of wetland plants in constructed wetlands receiving municipal effluent in tropical Australia. *Wat. Sci. Tech.* 35: 135-142.
- Griggs, G.B. and Hein, J.R. 1980. Source, dispersal, and clay mineral composition of fine-grained sediment off Central and Northern California. *Journal of Geology* 88: 541-556.
- Groisman, P.Y., T.R. Karl and R.W. Knight. 1994. Observed impact of snow cover on the heat balance and the rise of continental spring temperatures. *Science* 263: 198-200.
- Hamilton, W. J. 2004. “Tricolored Blackbird (*Agelaius tricolor*)” in The Riparian bird conservation plan: A strategy for reversing the decline of riparian-associated birds in California. California Partners in Flight. http://www.prbo.org/calpif/htmldocs/species/riparian/tricolored_blackbird.htm
- Hansen, S., H.E. Jensen, N.E. Nielsen, and H. Svendsen. 1990. DAISY: soil plan atmosphere system model. NPO–Research Report No. A10. The National Agency for Environmental Protection, Copenhagen, Denmark.
- Harder, J.D., and Kirkpatrick, R.L. 1994. “Physiological methods in wildlife research”, 275–306 in *Research and management techniques for wildlife and habitats*, 5th ed.; T.A. Bookhout, ed.; The Wildlife Society, Bethesda, MD.
- Harrington, J.M. 1996. California stream bioassessment procedures. California Department of Fish and Game, Water Pollution Control Laboratory, Rancho Cordova, CA.
- Harrington, J.M. 1999. An index of biological integrity for first to third order Russian River tributary streams. California Department of Fish and Game, Water Pollution Control Laboratory, Rancho Cordova, CA.
- Harris, L.D., T. Hctor, D. Maehr, and J. Sanderson. 1996. “The role of networks and corridors enhancing the value and protection of parks and equivalent areas”, 173-197 in *National Parks and Protected Areas*, R.G. Wright, ed.; Cambridge, England: Blackwell Science.
- Hayden, B.P. 1999. Climate change and extratropical storminess in the United States: An assessment. *Journal of the American Water Resources Association* 35(6): 1387-1398.
- Hennessy, K.J., J.M. Gregory, and J.F.B. Mitchell. 1997. Changes in daily precipitation under enhanced greenhouse conditions. *Climate Dynamics* 13(9): 667–680.
- Hill, M.P., 1997. The potential for the biological control of the floating aquatic fern *Azolla filiculoides* Lamarck (red water fern/rooivaring) in South Africa. Report to the Water Research Commission by the Plant Protection Research Institute Agricultural Research Council. WRC Report No. KV 100/97.

- Hilty, J.A., W.Z. Lidicker, Jr., and A.M. Merenlender. 2006. Corridor ecology: The science and practice of linking landscapes for biodiversity conservation. Island Press, Washington DC, USA.
- Hitchcock, C.S., and K.I. Kelson. 1998. Assessment of seismogenic sources between the Rodgers Creek and San Andreas Faults, Northwest San Francisco Bay Region, Sonoma County, California. U.S.G.S., National Earthquake Hazards Reduction Program, Award 1434-HQ-97-GR-03153.
- Honton, J., and A. Warwick Sears. 2006. Enhancing and Caring for the Laguna. Santa Rosa: Laguna Foundation.
- Hooke, R. LeB. 2002. Toward a uniform theory of clastic sediment yield in the fluvial systems. *Geological Society of America Bulletin* 112: 1778-1786.
- Horvatić, Janja, Vesna Peršić, and Melita Mihaljević. 2006. Bioassay method in evaluation of trophic conditions and nutrient limitation in the Danube wetland waters (1388–1426 r. km). *Hydrobiologia* 563: 453-463.
- Howe, G.E., R. Gillis, and R.C. Mowbray. 1998. Effect of chemical synergy and larval stage on the toxicity of atrazine and alachlor to amphibian larvae. *Environmental Toxicology and Chemistry* 17: 519–525.
- Huber, G.E., and A.A. Steuter. 1984. Vegetation profile and grassland bird response to spring burning. *Prairie Naturalist* 16:55–61.
- Huber, W.C., and R.E. Dickson. 1988. Storm water management model version 4. Users manual. EPA/600/3-88/001a (NTIS PB88-236641/AS). U.S. Environmental Protection Agency, Athens, GA.
- Huffman, M.E. and C.F. Armstrong. 1980. *Geology for planning in Sonoma County, California*
- Janda, R.J. and K.M. Nolan. 1979. “Stream sediment discharge in Northwestern California” in *Guidebook for a field trip to observe natural and management-related erosion in Franciscan terrain of Northern California*. Geological Society of America, Cordilleran Section. IV, 1-27.
- Karl, T.R. and R.W. Knight. 1998. Secular trends of precipitation amount, frequency, and intensity in the United States. *Bulletin of the American Meteorological Society* 79(2): 231-241.
- Kim, J., T.K. Kim, R.W. Arritt and N.L. Miller. 2002. Impacts of increased atmospheric CO² on the hydroclimate of the western United States. *Journal of Climate* 15: 1926-1943.
- Kiparsky, M., and P.H. Gleick. 2003. *Climate change and California water resources: A survey and summary of the literature*. Pacific Institute for Studies in Development, Environment, and Security.

- Kotliar, N. B. 2007. Olive-sided Flycatcher (*Contopus cooperi*): A Technical Conservation Assessment. Fort Collins: U.S. Geological Survey. <http://www.fs.fed.us/r2/projects/scp/assessments/olivesidedflycatcher.pdf>
- Kreitinger, K., and T. Gardali. 2006. Bringing birds back: A guide to habitat enhancement in riparian and oak woodlands for the North Bay region. California Partners in Flight, PRBO Conservation Science. <http://www.prbo.org/calpif>
- Kunz, T.H., and R.A. Martin. 1982. *Plecotus townsendii*. Mammalian Species, 175: 1-6.
- Kushlan, J.A., Steinkamp M.J., Parsons K.C, et al. 2002. Waterbird conservation for the Americas: The North American waterbird conservation plan, version 1. Waterbird Conservation for the Americas, Washington DC.
- Laguna Advisory Committee. January 1988. Report to the City of Sebastopol.
- Laguna de Santa Rosa Foundation and Sonoma Land Trust. 2003. Resource atlas and protection plan.
- Laguna de Santa Rosa Foundation. 2007. Ludwigia control project: Year two report. <http://www.lagunafoundation.org>
- Lambert, S.J. 1995. The effect of enhanced greenhouse warming on winter cyclone frequencies and strengths. *Journal of Climate* 8(5): 1447–1452.
- Lokemoen, J.T. 1973. Waterfowl production on stock-watering ponds in the northern plains. *J Range Manage* 26 179–184.
- Lowe, R.L. 1996. “Periphyton patterns in lakes” in Stevenson, R. J., Bothwell, M. L. & Lowe, R. L., eds.; *Algal Ecology: Freshwater Benthic Ecosystems*. Academic Press, New York, 57–76.
- Ludke, J.L., E.F. Hill, and M.P. Dieter. 1975. Cholinesterase (ChE) response and related mortality among birds fed ChE inhibitors. *Archives of Environmental Contamination and Toxicology* 3: 1-21.
- Mack, R.N., D. Simberloff, W.M. Lonsdale, H. Evans, M. Clout, and F.A. Bazzaz. 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689-710.
- Mao, K. 1992. How to select a computer model for storm water management. *Pollution Engineering*, 60-64.
- Mayer, P.M., S.K. Reynolds, M.D. McCutchen, and T.J. Canfield. 2006. Riparian buffer width, vegetative cover, and nitrogen removal effectiveness: A review of current science and regulations. EPA/600/R-05/118. U.S. Environmental Protection Agency. Cincinnati, OH.
- McCormick, P.V., and R.J. Stevenson. 1998. Periphyton as a tool for ecological assessment and management in the Florida Everglades. *Journal of Phycology* 34: 726–733.

- McKeon, T.J. and J.J. Segna. 1987. Selection criteria for mathematical models used in exposure assessments: surface water models. US EPA Office of Health and Environmental Assessment. Washington, D.C. EPA/600/8-87/042.
- Merritt, W.S., R.A. Letcher, and A.J. Jakeman. 2003. A review of erosion and sediment transport models. *Environmental Modeling and Software* 18: 761-769.
- Milliman, J.D and J.P.M. Syvitski. 1992. Geomorphic/Tectonic control of sediment discharge to the ocean: The importance of small mountainous rivers. *The Journal of Geology* 100:525-544.
- Montgomery, D.R. and W.E. Dietrich. 1994. "Landscape dissection and drainage area-slope thresholds", 221-246 in *Process models and theoretical geomorphology*, M.J. Kirkby ed., John Wiley and Sons.
- Morrison, M. L. 1986. "Bird populations as indicators of environmental change" 429-451 in Richard F. Johnston, ed., *Current Ornithology*, Volume 3. Plenum Press, New York.
- National Invasive Species Council. 2001. Meeting the invasive species challenge: National invasive species management plan.
- National Research Council. 1995. Review of EPA's environmental monitoring and assessment program: overall evaluation. National Research Council, National Academy Press, Washington, DC.
- Nature Conservancy. 2005. Indicators of hydrologic alteration user's manual, Version 7. The Nature Conservancy with Smythe Scientific Software and Totten Software Design.
- NatureServe. 2006. NatureServe Explorer: An online encyclopedia of life. Arlington, Virginia. <http://www.natureserve.org/explorer/servlet/NatureServe?searchName=Rana+boylei>
- Neitsch S.L., J.G. Arnold, J.R. Kiniry, L.R. Williams, and K.W. King. 2002. Soil and water assessment tool (SWAT) theoretical documentation.
- Noda, A. and T. Tokioka. 1989. The effect of doubling the CO² concentration on convective and non-convective precipitation in a general circulation model coupled with a simple mixed layer ocean model. *Journal of the Meteorological Society of Japan* 67: 1057-1069.
- Novotny, V. and H. Olem. 1994. *Water quality: prevention, identification, and management of diffuse pollution*. New York: Van Nostrand Reinhold.
- Nurnberg, G.K. 1984. The prediction of internal phosphorus load in lakes with anoxic hypolimnia. *Limnology and Oceanography*, 29: 111-124.
- Ohlendorf, H.M., R.L. Hothem, C.M. Bunck, and K.C. Marois 1990. Bioaccumulation of selenium in birds at Kesterson Reservoir, California. *Contam. Toxicol.* 19, 495-507.
- Olden, J. D. and N. L. Poff. 2003. Redundancy and the choice of hydrologic indices for characterizing streamflow regimes. *River Research and Applications* 19: 101-121.

- Pacific Southwest Inter-Agency Committee (PSIAC). 1968. Report of the water management subcommittee: factors affecting sediment yield and measures for the reduction of erosion and sediment yield.
- Pearson S.F., and D.A. Manuwal. 2001. Breeding bird response to riparian buffer width in managed Pacific Northwest douglas-fir forests. *Ecological Applications* 11: 840-853.
- Perrin, C., C. Michel, and V. Andreassian. 2001. Does a large number of parameters enhance model performance? Comparative assessment of common catchment model structures on 420 catchments. *Journal of Hydrology* 242, 275-301.
- Philip Williams and Associates, Ltd. (PWA). 2001. Laguna de Santa Rosa feasibility study: year one geomorphic investigation. PWA Project Number 1411-08.
- Philip Williams and Associates, Ltd. (PWA). 2004. Sediment sources, rate and fate in the Laguna de Santa Rosa, Volume II. PWA Project Number 1411-08.
- Poff, N.L. and J.V. Ward. 1989. Implications of streamflow variability and predictability for lotic community structure: a regional analysis of streamflow pattern. *Canadian Journal of Fisheries and Aquatic Sciences* 46: 1805-1818.
- Fehring, K. and T. Gardali. 2005. Bird inventory and monitoring at the Laguna de Santa Rosa. PRBO Conservatin Science Contribution #1332.
- Refsgaard, J.C. and B. Storm. 1995. "MIKE SHE", 809-846 in V.P. Singh, ed., *Computer models of watershed hydrology*. Water Resources Publications, Highlands Ranch, Colorado.
- Regan, C. 1996. Florida Bay's osprey, eagles and terrapin give blood. Everglades National Park: National Park Service. <http://www.nps.gov/archive/ever/current/fbn96a-7.htm>
- Rehn, A.C., and P.R. Ode. September 2005. Development of a bethic index of biotic integrity (B-IBI) for wadeable streams in northern coastal California and its application to regional 305(b) assessment.
- Reichholf, J. 1976. "The influence of recreation activities on waterfowl", 364-369 in *Proceedings of the international conference on conservation of wetlands and waterfowl*, M. Smart, ed.; Slimbridge (Glos) England: International Waterfowl Research Bureau.
- Reynolds, K.M. 2002. Ecosystem management decision support 3.0. General technical report. PNW-GTR-470. U.S. Forest Service, Pacific Northwest Research Station, Portland, OR. <http://www.fsl.orst.edu/emds/>.
- Reynolds, K.M. and P. F. Hessburg. 2005. Decision support for integrated landscape evaluation and restoration planning. *Forest Ecology and Management* 207: 263-278.
- Reynolds, K.M., M. Jensen, J. Andreason, and I. Goodman. 2000. Knowledge-based assessment of watershed condition. *Computers and Electronics in Agriculture*. 27: 315-333.

- Reynolds, K.M., P. Cunningham, L. Bednar, M. Saunders, M. Foster, R. Olson, D. Schmoldt, D. Latham, B. Miller, and J. Steffenson. 1996. A design framework for a knowledge-based information management system for watershed analysis in the Pacific Northwest U.S. *AI Applications* 10:9-22.
- Richter, B.D., J.B. Baumgartner, J. Powell, and D. P. Braun. 1996. A method for assessing hydrologic alteration within ecosystems. *Conservation Biology* 10: 1163-1174.
- Ritter, J.R. and W.M. Brown III. 1971. Turbidity and suspended sediment transport in the Russian River Basin, California. U.S. Geological Survey Open File Report.
- Salop, P., K. Abu-Saba, A. Gunther, A. Feng. 2002. Analysis of the Laguna creek watershed monitoring program, year 1, 2001-2002. City of Fremont. http://www.cleanwaterprogram.org/uploads/ACSed00-01_summary_final.pdf.
- Scheuhammer, A.M. 1987. The chronic toxicity of aluminium, cadmium, mercury, and lead in birds: A review. *Environ. Pollut.* 46(4): 263-95.
- Semlitsch, R. and J.R. Bodie. 2006. Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology* 17: 1219-1228.
- Shelton, A.C. January 1911. Nesting of the California Cuckoo. *The Condor*, XIII: 19-22.
- Smith, R.D. August 2006. Russian River watershed management plan: Synthesis report for baseline watershed assessment. U.S. Army Engineer Research and Development Center—Environmental Laboratory, Vicksburg, MS.
- Snyder, M.A., J.L. Bell, L.C. Sloan, P.B. Duffy and B. Govindasamy. 2002. Climate responses to a doubling of atmospheric carbon dioxide for a climatically vulnerable region. *Geophysical Research Letters* 29(11): 10.1029/2001GL014431.
- Sonoma County Flood Control and Water Conservation District (SCFCWCD). 1965. Flood!! December 1964-January 1965, Santa Rosa.
- Sonoma County Water Agency (SCWA). 2006. Fisheries enhancement program. http://www.scwa.ca.gov/environment/natural_resources/fisheries_enhancement.php
- Sonoma County Water Agency (SCWA). December 2006. 2005 Urban water management plan.
- Steefel, C.I., Van Cappellan, P. 1998. Reactive transport modeling of natural systems. *Journal of Hydrology* 209, 1-7.
- Sustainable Land Stewardship Institute. 2005. Biological and physical/habitat assessment of selected sites within the city of Santa Rosa.
- Suttle, K.B., M.A. Thomsen, and M.E. Power. 2007. Species interactions reverse grassland responses to changing climate. *Science* 315: 640-642.
- Temple, S. and J. Wiens. 1989. Bird populations and environmental change: Can birds be bioindicators. *American Birds* 43: 260-270.

- Tetra Tech. 2006. Technical approach to develop nutrient numeric endpoints for California. U.S. EPA Region IX (Contract No. 68-C-02-108). Lafayette, CA.
- Todd Engineers. June 2004. Groundwater study for the Canon Manor West subdivision assessment district.
- Trso, M. October 10, 2006. Final sediment source analysis: Sonoma creek watershed, California. Sonoma Ecology Center with Watershed Sciences.
- US Army Corps of Engineers (USACE), San Francisco District. 2002. Santa Rosa Creek basin hydrology assessment.
- US Army Corps of Engineers (USACE), San Francisco District. 2003. Laguna de Santa Rosa feasibility study basin hydrology assessment.
- US Army Corps of Engineers (USACE), San Francisco District. 2004. Biological assessment for agency activities that may affect protected species of steelhead trout, coho salmon, and chinook salmon in the Russian River. http://www.spn.usace.army.mil/ets/rrsection7/Sec%206%20Final%20BA_integration_effects.pdf
- US Environmental Protection Agency (USEPA). September 1995. Bioindicators for assessing ecological integrity of prairie wetlands. Report # EPA/ 600/ R-96/ 082. http://www.epa.gov/owow/wetlands/wqual/pph2_6.html
- US Environmental Protection Agency (USEPA). 1997. Compendium of tools for watershed assessment and TMDL development. EPA841-B-97-006.
- US Environmental Protection Agency (USEPA). 1999. Protocol for developing nutrient TMDLs. EPA 841-B-99-007. Washington DC.
- US Environmental Protection Agency (USEPA). 2006. Freshwater fish identification and their use as indicators. <http://www.epa.gov/bioiweb1/html/fish.html>.
- US Fish and Wildlife Service (USFWS). May 23, 1996. Endangered and threatened wildlife and plants; determination of threatened status for the California red-legged frog. Federal Register, Vol. 61, No. 101. US Fish and Wildlife Service. http://ecos.fws.gov/docs/federal_register/fr2960.pdf.
- US Fish and Wildlife Service (USFWS). October 22, 1997. Endangered and threatened wildlife and plants; determination of endangered status for nine plants from the grasslands or mesic areas of the Central Coast of California. Federal Register, Volume 62, Number 204. US Fish and Wildlife Service. <http://www.cdpr.ca.gov/docs/es/estext/fr102297.txt>.
- US Fish and Wildlife Service (USFWS). 1998. Recovery plan for the California freshwater shrimp (*Syncaris pacifica* Holmes 1895). US Fish and Wildlife Service. http://ecos.fws.gov/docs/recovery_plans/1998/980731a.pdf.

- US Fish and Wildlife Service (USFWS). July 25, 2001. Endangered and threatened wildlife and plants; 12-month finding for a petition to list the yellow-billed cuckoo (*Coccyzus americanus*) in the western continental United States. Federal Register, Volume 66, Number 143. US Fish and Wildlife Service. http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=2001_register&docid=01-18560-filed.
- US Fish and Wildlife Service (USFWS). December 3, 2003. Notice of availability of the post-delisting monitoring plan for the American peregrine falcon (*Falco peregrinus anatum*). Federal Register, Volume 68, Number 232. US Fish and Wildlife Service. <http://www.fws.gov/policy/library/03-30065.html>.
- Weller, M. W., C.S. Spatcher. 1965. Role of habitat in the distribution and abundance of marsh birds. Iowa State University.
- Welsh, H. Jr., and L.M. Ollivier. 1998. Stream amphibians as indicators of ecosystem stress: A case study from California's redwoods. *Ecological Applications* 8: 118-1132.
- Weseloh, D.V.C., J. Rodrigue, H. Blokpoel, and P.J. Ewins. 1997. Contaminant concentrations in eggs of black terns (*Chlidonias niger*) from southern Ontario and southern Quebec, 1989-1996. *Colonial Waterbirds* 20: 604-616.
- Wickham, D. 2000. Phosphate loading and eutrophication in the Laguna de Santa Rosa. Report to Russian River watershed protection committee and City of Santa Rosa.
- Wilcock, P. et al. 2003. "When models meet managers: Examples from geomorphology" in *Prediction in Geomorphology*. American Geophysical Union. <http://www.fsl.orst.edu/wpg/pubs/03Wilcocketal.pdf>.
- Williams, J.R., March 7-9, 1981. Testing the modified universal soil loss equation (MUSLE), 157-165 in *Proceedings of the workshop on estimating erosion and sediment yield on rangelands*. Tucson, Arizona. USDA Agricultural Research Service, Agricultural Reviews and Manuals ARM-W-26.
- Winzler and Kelly, and Luhdorff and Scalmanini. January 2005. City of Rohnert Park water supply assessment.
- Winzler and Kelly. March 2004. Draft southern Santa Rosa area drainage study. City of Santa Rosa.
- Wohl, E., R. McConnell, J. Skinner, and R. Stenzel. 1998. Inheriting our past: River sediment sources and sediment hazards in Colorado. *Water in the balance report No.7*. Colorado Water Resources Research Institute, Colorado State University, Fort Collins, Colorado.

